

WUE-ITP-97-036
hep-ph/9709345

August 1997

The Parton Structure of Real Photons^{*}

Andreas Vogt

*Institut für Theoretische Physik, Universität Würzburg
Am Hubland, D-97074 Würzburg, Germany*

Abstract

The QCD treatment of the photon structure is recalled. Emphasis is given to the recently derived momentum sum rule, and to the proper choice of the factorization scheme and/or boundary conditions for the evolution equations beyond the leading order. Parametrizations of the photon's parton content are examined and compared. The small- x behaviour of the photon structure is briefly discussed.

^{*} Invited talk presented at PHOTON '97, Egmond aan Zee, The Netherlands, May 1997. To appear in the proceedings.

THE PARTON STRUCTURE OF REAL PHOTONS

ANDREAS VOGT

*Institut für Theoretische Physik, Universität Würzburg,
Am Hubland, D-97074 Würzburg, Germany*

The QCD treatment of the photon structure is recalled. Emphasis is given to the recently derived momentum sum rule, and to the proper choice of the factorization scheme and/or the boundary conditions for the evolution equations beyond the leading order. Parametrizations of the photon's parton content are examined and compared. The small- x behaviour of the photon structure is briefly discussed.

1 Introduction

Deep-inelastic electron-photon scattering has been the classical process for investigating the hadronic structure of the photon¹. This process is kinematically analogous to the usual lepton-nucleon scattering. It has quite early received special interest, since the structure function $F_2^\gamma(x, Q^2)$ can be completely calculated in perturbation theory^{2,3} at large Bjorken- x and large resolution Q^2 . At scales accessible at present or in the near future, however, these results are unfortunately not applicable. Hence the photon structure functions have to be analyzed in terms of non-perturbative initial distributions for the QCD evolution equations⁴, very much like the nucleon case.

Experimentally F_2^γ has been determined, albeit with rather limited accuracy, via $e^+e^- \rightarrow e^+e^- + \text{hadrons}$ at all electron-positron colliders since PEP and PETRA. The longitudinal structure function F_L^γ has been inaccessible so far, and will presumably remain so in the foreseeable future^{5,6}. On the other hand, the past months have witnessed a substantial amount of new results on F_2^γ from LEP, and many more can be expected from forthcoming LEP2 runs. If systematic problems in extractions of F_2^γ from final-state modeling⁶ can be overcome, these results will be able to challenge seriously the present, model-driven theoretical understanding of the photon structure.

In this talk a brief survey is given of the present theoretical and phenomenological status of this subject. In Section 2 we recall the evolution equations for the photon's parton distributions, including the recently derived momentum sum rule. The factorization scheme ambiguities are more relevant here as in the usual hadronic case, this issue is discussed in Section 3. Some of the most relevant parametrizations of the quark and gluon densities of the photon are discussed in Section 4 with respect to their assumptions and limitations. Finally Section 5 is devoted to the small- x behaviour of the photon structure functions. For other aspects the reader is referred to refs.¹.

2 The evolution of the photon's parton densities

The photon is a genuine elementary particle, unlike the hadrons. Hence it can directly take part in hard scattering processes, in addition to its quark and gluon distributions arising from quantum fluctuations, $q^\gamma(x, Q^2)$ and $g^\gamma(x, Q^2)$. Denoting the corresponding photon distribution in the photon by $\Gamma^\gamma(x, Q^2)$, the evolution equations for these parton densities are generally given by

$$\begin{aligned}\frac{dq_i^\gamma}{d\ln Q^2} &= \frac{\alpha}{2\pi} \bar{P}_{q_i\gamma} \otimes \Gamma^\gamma + \frac{\alpha_s}{2\pi} \left\{ 2 \sum_{k=1}^f \bar{P}_{q_i q_k} \otimes q_k^\gamma + \bar{P}_{q_i g} \otimes g^\gamma \right\} \\ \frac{dg^\gamma}{d\ln Q^2} &= \frac{\alpha}{2\pi} \bar{P}_{g\gamma} \otimes \Gamma^\gamma + \frac{\alpha_s}{2\pi} \left\{ 2 \sum_{k=1}^f \bar{P}_{g q_k} \otimes q_k^\gamma + \bar{P}_{g g} \otimes g^\gamma \right\} \\ \frac{d\Gamma^\gamma}{d\ln Q^2} &= \frac{\alpha}{2\pi} \bar{P}_{\gamma\gamma} \otimes \Gamma^\gamma + \frac{\alpha_s}{2\pi} \left\{ 2 \sum_{k=1}^f \bar{P}_{\gamma q_k} \otimes q_k^\gamma + \bar{P}_{\gamma g} \otimes g^\gamma \right\}.\end{aligned}\quad (1)$$

Here $\alpha \simeq 1/137$ is the electromagnetic coupling constant, and $\alpha_s \equiv \alpha_s(Q^2)$ denotes the running QCD coupling. $\otimes$ represents the Mellin convolution, and f stands for the number of active (massless) quark flavours. The antiquark distributions do not occur separately in Eq. (1), as $\bar{q}_i^\gamma(x, Q^2) = q_i^\gamma(x, Q^2)$ due to charge conjugation invariance. The generalized splitting functions read

$$\bar{P}_{ij}(x, \alpha, \alpha_s) = \sum_{l,m=0} \frac{\alpha^l \alpha_s^m}{(2\pi)^{l+m}} \bar{P}_{ij}^{(l,m)}(x), \quad (2)$$

with $\bar{P}_{q_i q_k}$ being the average of the quark–quark and antiquark–quark splitting functions. The parton densities are subject to the energy–momentum sum rule

$$\int_0^1 dx x \left[\Sigma^\gamma(x, Q^2) + g^\gamma(x, Q^2) + \Gamma^\gamma(x, Q^2) \right] = 1, \quad (3)$$

where Σ represents the singlet quark distribution, $\Sigma^\gamma = 2 \sum_{i=1}^f q_i^\gamma$.

Usually calculations involving the photon's parton structure are restricted to first order in $\alpha \ll 1$. In this approximation all $l \neq 0$ terms in Eq. (2) can be neglected, since q_i^γ and g^γ are already of order α . This reduces the functions $\bar{P}_{ij}$ to the usual QCD quantities $P_{ij}(x, \alpha_s)$, with $P_{\gamma q_i}$ and $P_{\gamma g}$ dropping out completely. Moreover one has $P_{\gamma\gamma} \propto \delta(1-x)$ to all orders in α_s , as real photon radiations from photons starts at order α^2 only. Thus the last line of Eq. (1) can be integrated immediately, at leading order (LO), $m = 0$, resulting in

$$\Gamma_{\text{LO}}^\gamma(x, Q^2) = \delta(1-x) \left[1 - \frac{\alpha}{\pi} \left(\sum_q e_q^2 \ln \frac{Q^2}{Q_0^2} + c_1 \right) \right]. \quad (4)$$

Here e_q stands for the quark charges, Q_0^2 is some reference scale for the evolution, and the constant c_1 will be discussed below. Only the $O(1)$ part of Γ^γ affects the quark and gluon densities at order α , as well as any observable involving hadronic final states like F_2^γ , leading to the evolution equations⁷

$$\begin{aligned}\frac{dq_i^\gamma}{d\ln Q^2} &= \frac{\alpha}{2\pi}P_{q_i\gamma} + \frac{\alpha_s}{2\pi}\left\{2\sum_{k=1}^f P_{q_i q_k} \otimes q_k^\gamma + P_{q_i g} \otimes g^\gamma\right\} \\ \frac{dg^\gamma}{d\ln Q^2} &= \frac{\alpha}{2\pi}P_{g\gamma} + \frac{\alpha_s}{2\pi}\left\{2\sum_{k=1}^f P_{g q_k} \otimes q_k^\gamma + P_{gg} \otimes g^\gamma\right\}.\end{aligned}\quad (5)$$

The splitting functions $P_{ij}(x, \alpha_s)$ are presently known to next-to-leading order (NLO) in α_s , $m = 1$, see refs.^{8,9,10}.

The momentum sum rule (3) holds order by order in α , thus Eq. (4) implies

$$\int_0^1 dx x \left[\Sigma_{\text{LO}}^\gamma(x, Q^2) + g_{\text{LO}}^\gamma(x, Q^2) \right] = \frac{\alpha}{\pi} \left(\sum_q e_q^2 \ln \frac{Q^2}{Q_0^2} + c_1 \right). \quad (6)$$

The photon's quark and gluon densities are therefore not related by a hadron-type sum rule. Instead their momentum fractions rise logarithmically with Q^2 as long as the lowest-order approximation in α is justified. Hence, on the level of Eq. (5) alone, an important constraint on the parton densities is missing. That deficit can be removed by inferring c_1 from elsewhere, as recently done in refs.^{11,12} by connecting Eq. (4) to the cross section $\sigma(e^+e^- \rightarrow \text{hadrons})$ via a dispersion relation in the photon virtuality. This procedure yields¹¹

$$\left(\frac{c_1}{\pi}\right)_{\text{LO}} = \sum_{V=\rho,\omega,\phi} \frac{4\pi}{f_V^2} \simeq 0.55 \quad \text{at} \quad Q_0^2 \simeq (0.6 \text{ GeV})^2. \quad (7)$$

An error of about 20% can be assigned to this value, arising from the uncertainties of f_ρ^2 (leptonic ρ width vs. $\gamma p \rightarrow \rho^0 p$) and of the scale Q_0^2 where the connection of c_1 to the vector-meson decay constants holds. The numerical results of refs.^{11,12} agree well within this margin.

The general solution of the inhomogeneous evolution equations (5) reads

$$\vec{q}^\gamma = \begin{pmatrix} \Sigma^\gamma \\ g^\gamma \end{pmatrix} = \vec{q}_{\text{PL}}^\gamma + \vec{q}_{\text{had}}^\gamma, \quad (8)$$

where only the flavour singlet part has been indicated. The solution of the homogeneous ('hadronic') equation, $\vec{q}_{\text{had}}^\gamma(x, Q^2)$, contains the perturbatively uncalculable boundary conditions $\vec{q}^\gamma(x, Q_0^2)$. The inhomogeneous ('pointlike') part, on the other hand, is completely calculable once Q_0^2 has been specified.

At next-to-leading order these solutions can be written as^{10,13}

$$\vec{q}_{\text{had}}^\gamma = \left(\left[\frac{\alpha_s}{\alpha_0} \right]^{\hat{d}} + \frac{\alpha_s}{2\pi} \left\{ \hat{U} \otimes \left[\frac{\alpha_s}{\alpha_0} \right]^{\hat{d}} - \left[\frac{\alpha_s}{\alpha_0} \right]^{\hat{d}} \otimes \hat{U} \right\} \right) \otimes \vec{q}^\gamma(Q_0^2) \quad (9)$$

and

$$\vec{q}_{\text{PL}}^\gamma = \left\{ \frac{2\pi}{\alpha_s} + \hat{U} \right\} \otimes \left\{ 1 - \left[\frac{\alpha_s}{\alpha_0} \right]^{1+\hat{d}} \right\} \otimes \frac{1}{1+\hat{d}} \otimes \vec{a} + \left\{ 1 - \left[\frac{\alpha_s}{\alpha_0} \right]^{\hat{d}} \right\} \otimes \frac{1}{\hat{d}} \otimes \vec{b} \quad (10)$$

with $\alpha_0 = \alpha_s(Q_0^2)$. $\vec{a}$, $\vec{b}$, $\hat{d}$ and $\hat{U}$ stand for known combinations of the splitting functions and the QCD β -function. The LO evolution is obtained from Eqs. (9) and (10) by putting $\hat{U} = 0$ and $\vec{b} = 0$. A convenient way to evaluate these expressions is by transformation to Mellin moments, which reduces the convolutions to simple products. The x -dependent distributions are then calculated by a numerical Mellin inversion of the final result (8).

3 Boundary conditions and factorization schemes

The structure function F_2^γ is, at first order in the electromagnetic coupling α ,

$$F_2^\gamma = \sum_{q=u,d,s} 2x e_q^2 \left\{ q^\gamma + \frac{\alpha_s}{2\pi} (C_{2,q} \otimes q^\gamma + C_{2,g} \otimes g^\gamma) + \frac{\alpha}{2\pi} e_q^2 C_{2,\gamma} \right\}. \quad (11)$$

Only the contribution of the light flavours has been written out here. The reader is referred to refs.^{14,15} for the heavy quark part $F_{2,h}^\gamma$. At LO in α_s , just the first term in Eq. (11) is taken into account since $q^\gamma \sim 1/\alpha_s$, see Eq. (10). At NLO the usual hadronic one-loop coefficient functions $C_{2,q}(x)$ and $C_{2,g}(x)$ enter, together with the direct-photon contribution $C_{2,\gamma}$ given by³

$$C_{2,\gamma}^{\overline{\text{MS}}}(x) = 3 \left([x^2 + (1-x^2)] \ln \frac{1-x}{x} - 1 + 8x(1-x) \right). \quad (12)$$

This term causes difficulties in this standard factorization scheme, as it leads to a large LO/NLO difference for the inhomogeneous part $F_{2,\text{PL}}^\gamma$. In particular it is strongly negative at large x , see Fig. 1. Thus $F_{2,\text{PL}}^\gamma$ turns positive over the full x -range, for $Q_0^2 = 1 \text{ GeV}^2$, only at $Q^2 \simeq 20 \text{ GeV}^2$. This unphysical behaviour has to be overcome in the complete F_2^γ by the $\overline{\text{MS}}$ initial distributions, which are therefore forced to be very different from their LO counterparts.

These problems are circumvented by adopting the DIS_γ scheme introduced in refs.^{10,16}. Here $C_{2,\gamma}$ is absorbed into the quark distributions according to

$$q_{\text{DIS}_\gamma}^\gamma = q_{\overline{\text{MS}}}^\gamma + \frac{\alpha}{2\pi} e_q^2 C_{2,\gamma}^{\overline{\text{MS}}}, \quad C_{2,\gamma}^{\text{DIS}_\gamma} = 0. \quad (13)$$




Figure 1: The pointlike structure function $F_{2,\text{PL}}^\gamma$ in LO and in NLO for the $\overline{\text{MS}}$ and DIS_γ schemes. $Q_0^2 = 1 \text{ GeV}^2$, three active flavours and $\Lambda_{\text{LO}} = \Lambda_{\text{NLO}} = 0.2 \text{ GeV}$ have been used.

The coefficient functions $C_{2,q}$ and $C_{2,g}$ in Eq. (11), as well as the definition of the gluon density remain unchanged, in contrast to the hadronic DIS scheme¹⁷. Therefore F_2^γ assumes the usual hadronic $\overline{\text{MS}}$ form without a direct term in DIS_γ , resulting in a good LO/NLO stability of $F_{2,\text{PL}}^\gamma$ as illustrated in Fig. 1. Consequently physically motivated boundary conditions for the quark and gluon densities can be employed in this scheme also beyond leading order. An additional advantage of the DIS_γ scheme is that the leading $\overline{\text{MS}}$ terms for $x \rightarrow 0$ cancel in the transformed NLO photon-parton splitting functions¹⁸

$$P_{q\gamma}^{(1)} \sim \begin{cases} \ln^2 x + \dots \\ 2 \ln x + \dots \end{cases}, \quad P_{g\gamma}^{(1)} \sim \begin{cases} 1/x + \dots & \overline{\text{MS}} \\ -3 \ln x + \dots & \text{DIS}_\gamma \end{cases}. \quad (14)$$

In fact, this cancellation of the leading small- x term of $P_{g\gamma}^{\overline{\text{MS}}}$ does not only take place at NLO, but persists to all orders in α_s ¹⁹.

An equivalent $\overline{\text{MS}}$ formulation of the above solution to the $C_{2,\gamma}$ problem has been pursued in refs^{20,21}. It can be written as a modification of the pointlike part ($\overline{\text{PL}}$) in Eq. (8) by an additional ‘technical’ NLO input density,

$$q_{\overline{\text{PL}}}^\gamma(x, Q_0^2) = -\frac{\alpha}{2\pi} e_q^2 C_{2,\gamma}^{\overline{\text{MS}}}(x), \quad g_{\overline{\text{PL}}}^\gamma(x, Q_0^2) = 0. \quad (15)$$

This leads to $F_{2,\overline{\text{PL}}}^\gamma(x, Q_0^2) = 0$ and thus allows for similar ‘physical’ initial distributions on top of Eq. (15). The resulting quark distributions, however,




Figure 2: The pointlike up-quark density u_{PL}^γ in LO and in NLO for the DIS_γ scheme, compared to the physically equivalent $\overline{\text{MS}}$ distribution ($\overline{\text{PL}}$) with the input (15). The parameters Q_0^2 , f and Λ are as in Fig. 1. Also shown is the result for the PL' boundary condition (16).

exhibit a rather unphysical shape. As displayed in Fig. 2, they are suppressed (strongly enhanced) at medium (large) values of x , respectively, with respect to the pointlike LO and DIS_γ results. In a fully consistent NLO calculation, this $\overline{\text{MS}}$ procedure is nevertheless strictly equivalent to the DIS_γ treatment. On the other hand, as soon as not all terms beyond NLO are carefully omitted, the $\overline{\text{MS}}$ treatment turns out to be unstable at large x , see refs. ^{22,23}.

Due to the non-universality of the coefficient function $C_{2,\gamma}$, a special role is assigned to F_2^γ in the redefinitions (13,15) of the quark densities, similar to the hadronic DIS scheme. An alternative process-independent approach was worked out in ref. ²⁴. A universal technical $\overline{\text{MS}}$ input has been inferred from a detailed analysis of the Feynman diagrams for $\gamma^*\gamma \rightarrow \gamma^*\gamma$, which leads to

$$q_{\text{PL}'}^\gamma(x, Q_0^2) = -\frac{\alpha}{2\pi} e_q^2 C_\gamma'(x), \quad g_{\text{PL}'}^\gamma(x, Q_0^2) = 0 \quad (16)$$

with

$$C_\gamma'(x) = 3 \left([x^2 + (1-x^2)] \ln(1-x) + 2x(1-x) \right). \quad (17)$$

The resulting modified pointlike structure function $F_{2,\text{PL}'}^\gamma$, also shown in Fig. 1, remains negative at large x due to the uncompensated -1 in Eq. (12) only close to the reference scale Q_0^2 . At medium to small x , $F_{2,\text{PL}'}^\gamma$ is similar to the pointlike $\overline{\text{MS}}$ results, i.e., larger than its LO and $\text{DIS}_\gamma = \overline{\text{PL}}$ counterparts. The corresponding up-quark distributions are also illustrated in Fig. 2.

4 Parametrizations of photonic parton distributions

In order to specify the photon's parton densities, the perturbatively uncalculable initial distributions, $q_i^\gamma(x, Q_0^2)$ and $g^\gamma(x, Q_0^2)$, have to be fixed at some scale Q_0^2 . Only one combination of quark densities (dominated by u^γ) is presently well constrained, however, by F_2^γ data at $0.01 \lesssim x \lesssim 0.8$ from PETRA^{25,26,27}, PEP²⁸, TRISTAN^{29,30}, and LEP^{31,32,33}. The complete present data, including new results presented at this conference, is shown in Fig. 3 together with the NLO parametrizations of refs.^{16,24}. The gluon distribution is not tightly constricted either: there is sound evidence for $g^\gamma \neq 0$, and a very large and hard g^γ has been ruled out by jet production results^{34,35,36}.

Due to these limitations, current parametrizations invoke theoretical estimates and model assumptions, in particular from vector meson dominance (VMD). For safely high reference scales, $Q_0^2 \gtrsim 2 \text{ GeV}^2$, however, purely hadron-like initial distributions are known to be insufficient. An additional hard quark component has to be supplemented there in order to meet the F_2^γ data at larger Q^2 . In view of this situation two approaches have been used. First one can keep $Q_0 \geq 1 \text{ GeV}$, fit the quark densities to F_2^γ data, and estimate the gluon input. This method has been adopted in refs.^{37,38} and, more recently, in refs.^{11,21,39,40}. The second option is to retain a pure VMD ansatz,

$$(q_i^\gamma, g^\gamma)(x, Q_0^2) = \frac{4\pi\alpha}{f_\rho^2} (q_i^\rho, g^\rho)(x, Q_0^2) + \dots, \quad (18)$$

together with assumptions on the experimentally unknown ρ distributions, and to start the evolution at a very low scale $Q_0 \simeq 0.5 \dots 0.7 \text{ GeV}$ ^{11,16,24,41}. Note that this boundary condition complies with the momentum sum rule (7) if the ω and ϕ contributions are appropriately added.

In the following, the three available NLO parametrizations^{16,21,24} are briefly compared, together with the recent LO sets of ref.¹¹. For all these distributions $\Lambda_{\text{LO}, \overline{\text{MS}}} = 200 \text{ MeV}$ have been employed at $f = 4$.

The resulting u -quark densities are displayed in Fig. 4. Considering the LO results first, one notices that the parametrizations form two groups in the well-measured intermediate x -range, $0.2 \lesssim x \lesssim 0.7$. The lower one consists of the two low- Q_0 sets, GRV¹⁶ and SAS1D¹¹, which start the evolution at $Q_0^2 = 0.25 \text{ GeV}^2$ and 0.36 GeV^2 , respectively. The reference scales for the higher SAS2D^a, and GS(96)²¹ distributions read $Q_0^2 = 4 \text{ GeV}^2$ and 3 GeV^2 . This difference has been driven at least partly by first LEP data^{32,42}, see ref.⁴³, which were considerably higher than previous results around $x = 0.2$. A more consistent picture is now emerging from the new LEP data in this range.

^aThe additional SaS1M and SaS2M sets in ref.¹¹ are theoretically inconsistent, as the leading-order evolution is combined with the scheme-dependent coefficient function $C_{2,\gamma}$.




Figure 3: The presently available F_2^γ data compared to NLO parametrizations of refs. ^{16,24}. The hadron-like VMD components of the latter are separately displayed at $Q^2 = 2.4$ GeV².




Figure 4: Parametrizations of the up-quark distribution at LO^{11,16,21} and NLO^{16,21,24}. NLO results in $\overline{\text{MS}}$ have been transformed to the DIS_γ scheme according to Eq. (13).

The second striking feature in Fig. 4 is the large- x behaviour of the GS parametrization. Unlike SaS 2D, where a simple hard term $\propto x$ is employed, GS choose the massive Born expressions for $\gamma^*\gamma \rightarrow q\bar{q}$ at Q_0^2 on top of the hadronic VMD input. All power-law contributions $O([m_q^2/Q^2]^n)$, $n \geq 1$, are retained, resulting in a threshold at $x \simeq 0.9$ for typical constituent quark masses. Such a procedure, however, may be considered as inadequate for the construction of leading-twist parton densities.

Let us now turn to the NLO distributions. The results of GRV¹⁶ and AFG²⁴ are both based on the VMD ansatz (18), imposed at $Q_0^2 = 0.3 \text{ GeV}^2$ and 0.5 GeV^2 , respectively. The differences between these two parametrizations at $x \gtrsim 0.1$ can be understood in terms of the non-hadronic NLO boundary conditions discussed in Sec. 3, cf. Fig. 1. At lower x , the deviations are dominated by the differing assumptions^{44,45} on the experimentally virtually unconstrained pion sea – both groups estimate the unknown ρ distributions by their pionic counterparts. The third NLO set, GS(96), is technically flawed: it should at Q_0^2 , by construction, lead to the same F_2^γ results as the LO fit. However, u^γ turns out to be sizeably too small over the full x -range. Hence this parametrization is unfortunately not usable in its present form^b.

^bThis discussion also applies to the previous NLO parametrization²⁰ of the same group.




Figure 5: Parametrizations of the photon's gluon density at LO^{11,16,21} and NLO^{16,21,24}.

Before considering the gluon density, it is appropriate to comment on the quark-flavour decomposition and the momentum sum rule. Both SaS1D and AFG perform a coherent addition of the three light vector mesons at their respective input scales, with slightly differing assumptions on SU(3) breaking and the value of f_ρ . That leads to a suppression of the d -valence density by a factor of four with respect to the ρ -meson's u -valence component. This approach is able to describe the F_2^γ data without any further adjustment, hence the momentum sum rule (7) is met in both cases. On the other hand, GRV use just a ρ distribution, with a prefactor adjusted to the data. Although a factor of 1.6 perfectly mimics the F_2 of the (SU(3) symmetric) coherent superposition, too much momentum is spent due to the $u_v = d_v$ symmetry. Thus Eq. (6) is violated, e.g., by about +40% in LO at $Q^2 = 4 \text{ GeV}^2$. Finally the high- Q_0 fits, SaS2D and GS, do not impose the momentum sum rule at all.

The gluon distributions of these parametrizations are finally presented in Fig. 5. The pion distributions of AFG⁴⁴ and GRV⁴⁵ both describe the direct-photon production data in πp collisions⁴⁶, that is why these photonic gluons are so similar except at very x . For the GS parametrization, the gluon densities have been constrained by a LO comparison to TRISTAN jet production data^{34,35}, which seem to prefer a relatively large gluon distributions. The shapes of the SaS gluon densities are fixed by theoretical estimates, no direct or indirect experimental constraint has been imposed here.

5 Photon structure at small x

The region of very small parton momenta, $10^{-5} \lesssim x \lesssim 10^{-2}$, has attracted considerable interest in the proton case since the advent of HERA. The quark and gluon distributions show a marked rise at small x ^{47,48}, in good agreement with perturbative predictions for a low input scale $Q_0 \simeq 600$ MeV⁴⁹. The corresponding NLO evolution of the photon structure is shown in Fig. 6.




Figure 6: The NLO small- x evolution of the photon's quark and gluon distributions as predicted in ref¹⁶. The hadronic (VMD) and pointlike contributions are shown separately.

The parton distributions of the photon behave very differently in the limits of large and small x . In the former case, the perturbative part (10) dominates, especially for the quark distributions. On the other hand, this calculable contribution amounts at most to about 20% at very small- x , at scales accessible in the foreseeable future, in LO as well as in NLO. One may therefore expect, by VMD arguments for the hadronic component (9), a very similar rise as observed in the proton case. It will be very interesting to see whether this expectation is borne out by future F_2^γ measurements.

Let me finally mention that there is an even more intriguing, if exotic, possibility here: the calculable pointlike contribution could be drastically enhanced by large logarithmic small- x terms in the perturbation series¹⁹. If this component could be projected out, for example, by final-state observables, it would provide a rather unique small- x QCD laboratory.

Acknowledgement

This work has been supported by the German Federal Ministry for Research and Technology (BMBF) under contract No. 05 7WZ91P (0).

References

1. Ch. Berger and W. Wagner, *Phys. Rep.* **146**, 1 (1987);
H. Abramowicz *et al*, *Int. J. Mod. Phys.* **A8**, 1005 (1993);
M. Drees and R. Godbole, *J. Phys.* **G21**, 1559 (1995).
2. E. Witten, *Nucl. Phys.* **B120**, 189 (1977).
3. W.A. Bardeen and A.J. Buras, *Phys. Rev.* **D20** 166, (1979), E: **D21** 2041, (1980).
4. M. Glück and E. Reya, *Phys. Rev.* **D28**, 2749 (1983);
M. Glück, K. Grassie and E. Reya, *Phys. Rev.* **D30**, 1447 (1984).
5. P. Aurenche *et al*, in: *Proceedings of the LEP2 Physics Workshop*, eds. G. Altarelli, T. Sjöstrand and F. Zwirner (CERN 1996), p. 291.
6. D.J. Miller, **hep-ex/9708002**, these proceedings.
7. R.J. De Witt *et al*, *Phys. Rev.* **D19**, 2046 (1979), E: **D20**, 1751 (1979).
8. G. Curci, W. Furmanski and R. Petronzio, *Nucl. Phys.* **B175**, 27 (1980);
W. Furmanski and R. Petronzio, *Phys. Lett.* **B97**, 437 (1980);
E.G. Floratos, C. Kounnas and R. Lacaze, *Nucl. Phys.* **B192**, 417 (1981).
9. M. Fontannaz and E. Pilon, *Phys. Rev.* **D45**, 382 (1992), E: **D46**, 484 (1992).
10. M. Glück, E. Reya and A. Vogt, *Phys. Rev.* **D45**, 3986 (1992).
11. T. Sjöstrand and G.A. Schuler, *Z. Phys.* **C68**, 607 (1995);
G.A. Schuler in: *Proceedings of Photon '95*, eds. D.J. Miller, S.L. Cartwright and V. Khoze (World Scientific, Singapore, 1995).
12. L.L. Frankfurt and E.G. Gurvich, **hep-ph/9505406**; *J. Phys.* **G22**, 903 (1996).
13. W. Furmanski and R. Petronzio, *Z. Phys.* **C11**, 293 (1982).
14. E. Laenen, S. Riemersma, J. Smith and W.L. van Neerven, *Phys. Rev.* **D49**, 5753 (1994).
15. E. Laenen and S. Riemersma, in: *Proceedings of Photon '95*, eds. D.J. Miller, S.L. Cartwright and V. Khoze (World Scientific, Singapore, 1995);
Phys. Lett. **B376**, 169 (1996).
16. M. Glück, E. Reya and A. Vogt, *Phys. Rev.* **D46**, 1973, (1992).
17. G. Altarelli, R.K. Ellis and G. Martinelli, *Nucl. Phys.* **B157**, 461 (1979).
18. M. Glück, E. Reya and A. Vogt, *Phys. Rev.* **D48**, 116 (1993).
19. J. Blümlein and A. Vogt, DESY 96-096, August 1997.

20. L.E. Gordon and J.K. Storrow, *Z. Phys.* **C56**, 307 (1992).
21. L.E. Gordon and J.K. Storrow, *Nucl. Phys.* **B489**, 405 (1997).
22. M. Glück, E. Reya and A. Vogt, *Phys. Lett.* **B285**, 285 (1992).
23. A. Vogt in: *Proceedings of the Workshop on Two-Photon Physics at LEP and HERA*, eds. G. Jarlskog and L. Jönsson (Lund Univ., 1994), p. 141.
24. P. Aurenche, M. Fontannaz and J.P. Guillet, *Z. Phys.* **C64**, 621 (1994).
25. PLUTO Coll., Ch. Berger *et al*, *Phys. Lett.* **B142**, 111, (1984); *Nucl. Phys.* **B281**, 365 (1987).
26. JADE Coll., W. Bartel *et al*, *Z. Phys.* **C24**, 231 (1984).
27. TASSO Coll., M. Althoff *et al*, *Z. Phys.* **C31**, 527 (1986).
28. TPC/2 γ Coll., H. Aihara *et al*, *Phys. Rev. Lett.* **58**, 97 (1987); *Z. Phys.* **C34**, 1 (1987).
29. AMY Coll., S.K. Sahu *et al*, *Phys. Lett.* **B346**, 208 (1995); T. Kojima *et al*, *Phys. Lett.* **B400**, 395 (1997).
30. TOPAZ Coll., K. Muramatsu *et al*, *Phys. Lett.* **B332**, 477 (1994).
31. OPAL Coll., K. Ackerstaff *et al*, *Z. Phys.* **C74**, 33 (1997); CERN-PPE/97-087, CERN-PPE/97-104, July 1997; R. Nisius, these proceedings; J. Bechtluft, these proceedings.
32. DEPLHI Coll., P. Abreu *et al*, *Z. Phys.* **C69**, 223 (1996); I. Tyapkin, these proceedings; F. Kapusta *et al*, HEP'97, contribution 416.
33. ALEPH Coll., A. Finch *et al*, these proceedings; HEP'97, contribution 607.
34. TOPAZ Coll., H. Hayashij *et al*, *Phys. Lett.* **B314**, 149 (1993).
35. AMY Coll., B.J. Kim *et al*, *Phys. Lett.* **B325**, 248 (1994).
36. H1 Coll., T. Ahmed *et al*, *Nucl. Phys.* **B445**, 195 (1995).
37. M. Drees and K. Grassie, *Z. Phys.* **C28**, 451 (1985).
38. H. Abramowicz, K. Charchula and A. Levy, *Phys. Lett.* **B269**, 458 (1991).
39. K. Hagiwara, M. Tanaka, I. Watanabe, and T. Izubuchi, *Phys. Rev.* **D51**, 3197 (1995).
40. H. Abramowicz, E. Gurvich, and A. Levy, TAUP 2438-97, July 1997.
41. P. Aurenche *et al*, *Z. Phys.* **C56**, 589 (1992).
42. OPAL Coll., R. Akers *et al*, *Z. Phys.* **C61**, 199 (1994).
43. T. Sjöstrand, J.K. Storrow, and A. Vogt, *J. Phys.* **G22**, 893 (1996).
44. P. Aurenche *et al*, *Phys. Lett.* **B233**, 517 (1989).
45. M. Glück, E. Reya and A. Vogt, *Z. Phys.* **C53**, 651 (1992).
46. WA70 Coll., M. Bonesini *et al*, *Z. Phys.* **C37**, 535 (1988).
47. H1 Coll., S. Aid *et al*, *Nucl. Phys.* **B470**, 3 (1996).
48. ZEUS Coll., M. Derrick *et al*, *Z. Phys.* **C72**, 399 (1996).
49. M. Glück, E. Reya and A. Vogt, *Z. Phys.* **C53**, 127 (1992); **C67**, 433 (1995).